Minnesota – Rural Site

Total living turf cover [Total Cover (TC)] for Fall 2017 and Spring 2018 and percent total cover change from 2017 to 2018 (2017-2018 % TC Change). When treatment F tests were significant (p≤0.05), the Fisher's Protected Least Significant Difference Test (α=0.05) was used to separate means. For % TC Change (when significant), the most positive value within a column is shaded blue, while the most negative value within a column is shaded red. Remaining cells are shaded in a blue-purple-red gradient with zero shaded purple. Negative values indicate a reduction in turf coverage of desired species, whereas a positive value means the turf coverage of the plot increased.

This site was seeded 8/24/2017. The soil type was Cordova loam with 3.4% organic matter. This site was damaged due to a construction crane and had to be re-seeded in fall 2017. Plots were watered with a water truck during establishment.

		Fall 2017	Spring 2018	2017-2018
Turfgrass Entry	Species ¹	Total Cover (TC)		% TC Change
BAR PD06N17	AL	76.1	46.7	-29.4
BAR PD9032	AL	65.0	25.6	-39.4
Fults	AL	30.6	20.0	-10.6
Salton Sea	AL	61.7	11.7	-50.0
SeaSalt	AL	26.7	12.2	-14.4
Oceania Maritima	AM	63.9	41.1	-22.8
Castle	СН	78.9	8.9	-70.0
Compass II	СН	67.2	13.9	-53.3
FRC 43 M2	СН	81.1	8.3	-72.8
Heathland	СН	70.0	13.9	-56.1
Beacon	HD	92.2	28.9	-63.3
Gladiator	HD	84.4	25.6	-58.9
Nanook	HD	51.1	17.2	-33.9
Soil Guard	HD	71.7	7.2	-64.4
Sword	HD	77.2	12.2	-65.0
Barduke	КВ	56.7	7.8	-48.9
J-525	КВ	52.8	4.4	-48.3
J-793	КВ	37.8	2.2	-35.6
J-920	КВ	27.2	2.2	-25.0
Milagro	KB	57.2	1.1	-56.1
Morocco	KB	69.4	8.3	-61.1
Tirem	КВ	56.1	11.7	-44.4
Volt	КВ	72.8	0.6	-72.2
16-14-Lp 145	PR	90.6	1.1	-89.4
BAR Lp 10970	PR	77.8	2.8	-75.0
Gray Fox	PR	83.9	0.6	-83.3
Premium	PR	88.3	0.6	-87.8
Replicator ²	PR	90.6	1.7	-88.9
Stellar	PR	93.9	0.6	-93.3
BAR BIF 1GRL	SB	40.6	1.1	-39.4

Blue Mesa	SH	82.2	35.6	-46.7
J-248	SH	68.9	18.9	-50.0
Quatro	SH	81.7	23.9	-57.8
10RT DE	SL	90.0	29.4	-60.6
Sea Mist	SL	91.1	25.6	-65.6
Seabreeze GT	SL	84.4	12.8	-71.7
Shoreline	SL	87.8	37.2	-50.6
FRR 72 M2	ST	70.0	13.9	-56.1
Kent	ST	85.0	21.1	-63.9
Navigator II	ST	70.0	23.9	-46.1
Ruddy	ST	64.4	18.9	-45.6
Xeric	ST	81.1	18.9	-62.2
Avenger II	TF	87.2	3.9	-83.3
Birmingham	TF	86.1	10.6	-75.6
Black Tail	TF	68.9	5.0	-63.9
Fayette	TF	80.6	3.3	-77.2
JT-621	TF	73.9	1.1	-72.8
MNKY	TF	88.9	11.7	-77.2
Saltillo	TF	79.4	7.8	-71.7
Thunderstruck	TF	78.9	1.7	-77.2
MI DOT THV	MX	73.3	21.7	-51.7
MI DOT TUF	MX	71.1	34.4	-36.7
MN DOT 25-131	MX	83.9	31.7	-52.2
MN DOT MNST-12	MX	92.8	17.8	-75.0
NE DOT Rural Region 2	MX	81.7	9.4	-72.2
NE DOT Urban and Turf	MX	98.3	11.1	-87.2
NJ DOT A-4	MX	84.4	18.9	-65.6
NJ DOT Type B	MX	80.6	11.7	-68.9
WI DOT #20	MX	78.3	6.7	-71.7
WI DOT #40	MX	43.9	4.4	-39.4
LSD 0.05		30.8	18.6	27.4

¹Species tested include: AL = alkaligrass (*Puccinellia distans*); AM = seaside alkaligrass (*Puccinellia maritima*); CH = Chewings fescue (*Festuca rubra* ssp. *fallax*); HD = hard fescue (*Festuca brevipila*); KB = Kentucky bluegrass (*Poa pratensis*); PR = perennial ryegrass (*Lolium perenne*); SB = smooth bromegrass (*Bromus inermis*); SL = slender creeping red fescue (*Festuca rubra* ssp. *littoralis*); ST = strong creeping red fescue (*Festuca rubra* ssp. *rubra*); TF = tall fescue (*Schedonorus arundinaceus*); MX = DOT-specified mixtures. Mixture components can be found at http://roadsideturf.umn.edu/regional-roadside-state-mixtures.

²Tetraploid perennial ryegrass